arXiv:1106.2761v2 [math.DG] 22 Jun 2011

COMPACT CONFORMALLY KÄHLER EINSTEIN-WEYL MANIFOLDS

Włodzimierz Jelonek

Abstract. We give a description of compact conformally Kähler Einstein-Weyl manifolds whose Ricci tensor is Hermitian.

0. Introduction. In this paper we shall investigate compact Einstein-Weyl structures $(M,[g],D)$ on a complex manifold (M,J), $dim M\geq 4$, which are conformally Kähler and whose Ricci tensor ρ^D is Hermitian i.e. ρ^D is J -invariant.

We give a complete classification of compact Einstein-Weyl structures $(M,[g],D)$ with dim $M\geq 4$ such that $(M,[g],J)$ is conformally Kähler, i.e. there exists a metric $g_0\in[g]$ such that (M,g_0,J) is Kähler and whose Ricci tensor ρ^D is J-invariant i.e.

$$\rho^D(JX,JY)=\rho^D(X,Y).$$

Conformally Kähler Einstein manifolds were classified by A. Derdzinski and G. Maschler in [D-M-3]. The compact Einstein-Weyl 3-dimensional manifolds are studied in [T]. The compact Einstein-Weyl manifolds on complex manifolds compatible with complex structure are studied in [PS1], [PS2], [MPPS], [WW]. In [D-T] there are studied conditions on a Riemannian metric (M,g) in four dimensions for it to be locally conformal to Kähler. The Einstein-Weyl structures which are conformally Kähler studied in [MPPS], [WW] have the Ricci tensor ρ^D of the Weyl structure $(M,[g],D)$ which J -invariant.

1. Einstein-Weyl geometry and Killing tensors. We start with some basic facts concerning Einstein-Weyl geometry. For more details see [T], [PS1], [PS2].

Let M be a n-dimensional manifold with a conformal structure $[g]$ and a torsion-free affine connection D. This defines an Einstein-Weyl (E-W) structure if D preserves the conformal structure i.e. there exists a 1-form ω on M such that

$$Dg=\omega\otimes g \tag{1.1}$$

and the Ricci tensor ρ^D of D satisfies the condition

$$\rho^D(X,Y)+\rho^D(Y,X)=\bar{\Lambda}g(X,Y)\text{ for every }X,Y\in TM$$

for some function $\bar{\Lambda}\in C^\infty(M)$. P.Gauduchon proved ([G]) the fundamental theorem that if M is compact then there exists a Riemannian metric $g_0\in[g]$ for which $\delta\omega_0=0$ and g_0 is unique up to homothety. K.P. Tod proved [T] that the Gauduchon metric admits a Killing vector field dual to the form ω. We shall call g_0 a standard metric of E-W structure $(M,[g],D)$. Let ρ be a Ricci tensor of (M,g) and

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

let us denote by S the Ricci endomorphism of (M,g), i.e. $\rho(X,Y)=g(X,SY)$. We recall two important theorems (see [T], [PS1]):

Theorem 1.1. *A metric g and a 1-form ω define an E-W structure if and only if there exists a function $\Lambda\in C^\infty(M)$ such that*

$$\rho^\nabla+\frac{1}{4}\mathcal{D}\omega=\Lambda g \tag{1.2}$$

where $\mathcal{D}\omega=(\nabla_X\omega)Y+(\nabla_Y\omega)X+\omega(X)\omega(Y)$. If (1.2) holds then

$$\bar\Lambda=2\Lambda+div\omega-\frac{1}{2}(n-2)\parallel\omega^\sharp\parallel^2 \tag{1.3}$$

By ρ we shall denote the Ricci tensor of (M,g) and by $\tau=\operatorname{tr}_g\rho$ the scalar curvature of (M,g).

Definition. A Riemannian manifold (M,g) will be called a Gray $\mathcal{AC}^\perp$ manifold if the tensor $\rho-\frac{2\tau}{n+2}g$ is a Killing tensor.

In this paper Gray $\mathcal{AC}^\perp$ manifolds will be called for short Gray manifolds or $\mathcal{AC}^\perp$ manifolds.

Theorem 1.2. *Let M be a compact E-W manifold and let g be the standard metric with the corresponding 1-form ω. Then $\omega^\sharp$ is a Killing vector field on M.*

From the above theorems it follows (see [J])

Theorem 1.3. *Let $(M,[g])$ be a compact E-W manifold, $dimM\geq 3$, and let g be the standard metric on M. Then (M,g) is an $\mathcal{A}\oplus\mathcal{C}^\perp$-manifold. The manifold (M,g) is Einstein or the Ricci tensor ρ^∇ of (M,g) has exactly two eigenfunctions $\lambda_0\in C^\infty(M),\lambda_1=\Lambda$ satisfying the following conditions:*

(a) $(n-4)\lambda_1+2\lambda_0=C_0=const$

(b) $\lambda_0\leq\lambda_1$ on M

(c) $dimker(S-\lambda_0Id)=1, dimker(S-\lambda_1Id)=n-1$ on $U=\{x:\lambda_0(x)\neq\lambda_1(x)\}$.

In the addition $\lambda_0=\frac{1}{n}Scal_g^D$ where $Scal_g^D=tr_g\rho^D$ denotes the conformal scalar curvature of (M,g,D).

Note that $\omega(X)=g(\xi,X)$ where $\xi\in\mathfrak{iso}(M)$ and the formula

$$\rho^\nabla+\frac{1}{4}(n-2)\omega\otimes\omega=\Lambda g \tag{1.4}$$

holds. Thus $\nabla_X(\omega\otimes\omega)(X,X)=0$. From (1.4) it follows that

$$\nabla_X\rho(X,X)=X\Lambda g(X,X). \tag{1.5}$$

It means that $(M,g)\in\mathcal{A}\oplus\mathcal{C}^\perp$ and $d(\Lambda-\frac{2}{n+2}\tau)=0$, where τ is the scalar curvature of (M,g). From (1.5) it follows that the tensor $T=S-\Lambda Id$ is a Killing tensor. Let us denote by ξ the Killing vector field dual to ω. Note that $\rho(\xi,\xi)=(\Lambda-\frac{1}{4}(n-2)\parallel\xi\parallel^2)\parallel\xi\parallel^2$ and if $X\perp\xi$ then $SX=\Lambda X$. Hence the tensor S has two eigenfunctions $\lambda_0 = \Lambda - \frac{1}{4}(n-2) \parallel \xi \parallel^2$ and $\lambda_1 = \Lambda$. This proves (b). Note that

$$\tau = \lambda_0 + (n-1)\lambda_1 = n\Lambda - \frac{1}{4}(n-2) \parallel \xi \parallel^2 .$$

and $2\tau - (n+2)\Lambda = C_0 = const$. Thus $C_0 = (n-2)\Lambda - \frac{1}{2}(n-2) \parallel \xi \parallel^2$. However $(n-4)\lambda_1 + 2\lambda_0 = (n-2)\Lambda - \frac{1}{2}(n-2) \parallel \xi \parallel^2$ which proves (a). Note also that

$$\frac{1}{n} s_g^D = \Lambda - \frac{n-2}{4} \parallel \xi \parallel^2 = \lambda_0 \tag{3.6}$$

which finishes the proof.♢

On the other hand the following theorem holds (see [J]).

Theorem 1.4. *Let (M,g) be a compact $\mathcal{A} \oplus \mathcal{C}^{\perp}$ manifold. Let us assume that the Ricci tensor ρ of (M,g) has exactly two eigenfunctions λ_0, λ_1 satisfying the conditions:*

(a) $(n-4)\lambda_1 + 2\lambda_0 = C_0 = const$

(b) $\lambda_0 \leq \lambda_1$ on M

(c) $dimker(S - \lambda_0 Id) = 1, dimker(S - \lambda_1 Id) = n-1$ on $U = \{x : \lambda_0(x) \neq \lambda_1(x)\}$.

Then there exists a two-fold Riemannian covering (M', g') of (M,g) and a Killing vector field $\xi \in \mathfrak{iso}(M')$ such that $(M', [g'])$ admits two different E-W structures with the standard metric g' and the corresponding 1-forms $\omega_{\mp} = \mp \xi^{\sharp}$. The condition (b) may be replaced by the condition

(b1) there exists a point $x_0 \in M$ such that $\lambda_0(x_0) < \lambda_1(x_0)$.

Let us denote by τ the scalar curvature of (M,g). Then $\tau = (n-1)\lambda_1 + \lambda_0$ and $C_0 = (n-4)\lambda_1 + 2\lambda_0$. It follows that

$$\lambda_1 = \frac{2\tau - C_0}{n+2}, \quad \lambda_0 = \frac{(n-1)C_0 - (n-4)\tau}{n+2}. \tag{1.7}$$

In particular $\lambda_0, \lambda_1 \in C^{\infty}(M)$. Let S be the Ricci endomorphism of (M,g) and let us define the tensor $T := S - \lambda_1 Id$. Since from (1.7) we have $d\lambda_1 = \frac{2}{n+2} d\tau$ it follows that T is a Killing tensor with two eigenfunctions: $\mu = 0$ and $\lambda = \lambda_0 - \lambda_1$. Hence there exists a two fold Riemannian covering $p : (M', g') \to (M, g)$ and a Killing vector field $\xi \in \mathfrak{iso}(M')$ such that $S'\xi = (\lambda_0 \circ p)\xi$ where S' is the Ricci endomorphism of (M', g'). Note also that $\parallel \xi \parallel^2 = |\lambda - \mu| = |\lambda_0 - \lambda_1|$. Let us define the 1-form ω on M' by $\omega = c\xi^{\sharp}$ where $c = 2\sqrt{\frac{1}{n-2}}$. It is easy to check that with such a choice of ω equation (1.4) is satisfied and $\delta\omega = 0$. Thus (M', g', ω) defines an E-W structure and g' is the standard metric for $(M', [g'])$. Note that $(M, g', -\omega)$ gives another E-W structure corresponding to the field $-\xi$. ♢

Corollary 1.5. Let (M,g) be a compact simply connected manifold satisfying the assumptions of Th.3.4. Then $(M,[g])$ admits two E-W structures with the standard metric g.

2. Killing tensors. In this section we describe the Riemannian manifold (M,g) where $g \in [g])$ is the standard metric of Einstein-Weyl structure $(M, [g], D)$. We describe such manifolds under an additional condition, that the distribution corresponding to the eigenvalue λ_1 of the Ricci tensor is totally geodesic.

We say, that a distribution (not necessarily integrable) $\mathcal{D}$ is totally geodesic, if $\nabla_X X \in \Gamma(\mathcal{D})$ for every $X \in \Gamma(\mathcal{D})$.

We start with:

Lemma 2.1. *Let S be a self-adjoint tensor on (M,g) with exactly two eigenvalues λ,μ. If the distributions $\mathcal{D}_\lambda,\mathcal{D}_\mu$ are both umbilical, $\nabla\lambda \in \Gamma(D_\mu), \nabla\mu \in \Gamma(D_\lambda)$ and the mean curvatures ξ_λ,ξ_μ of the distributions D_λ, D_μ respectively satisfy the equations*

$$\xi_\lambda = \frac{1}{2(\mu-\lambda)}\nabla\lambda, \qquad \xi_\mu = \frac{1}{2(\lambda-\mu)}\nabla\mu,$$

then S is a Killing tensor.

Proposition 2.2. *Let (M,g) be a $2n$-dimensional Riemannian manifold whose Ricci tensor ρ has two eigenvalues $\lambda_0(x),\lambda_1(x)$ of multiplicity 1 and $2n-1$ respectively at every point x of M. Assume that the eigendistribution $\mathcal{D}_{\lambda_1}$ corresponding to λ_1 is totally geodesic. Then (M,g) is a Gray manifold if and only if $2\lambda_0+(n-4)\lambda_1$ is constant and $\nabla\tau \in \Gamma(\mathcal{D}_{\lambda_1})$.*

Proof. Let S_0 be the Ricci endomorphism of (M,g), i.e. $\rho(X,Y) = g(S_0X,Y)$. Let S be the tensor defined by the formula

$$S_0 = S + \frac{\tau}{n+1}\,\mathrm{id}\,. \tag{2.1}$$

Then

$$\operatorname{tr} S = -\frac{(n-1)\tau}{n+1}. \tag{2.2}$$

Let λ_0,λ_1 be the eigenfunctions of S_0 and let us assume that

$$2\lambda_0 + (2n-4)\lambda_1 = C \tag{2.3}$$

where $C \in \mathbb{R}$. Note that S also has two eigenfunctions which we denote by λ_0',λ_1' respectively. It is easy to see that $\lambda_0' = -\frac{n-1}{n+1}\tau + C\frac{2n-1}{2(n+1)}, \lambda_1' = -\frac{C}{2(n+1)}$ and $\lambda_0 = -\frac{\tau(n-2)}{n+1} + C\frac{2n-1}{2(n+1)}, \lambda_1 = \frac{\tau}{(n+1)} - \frac{C}{2(n+1)}$. Since the distribution $\mathcal{D}_{\lambda_0}$ is umbilical we have $\nabla_X X_{|\mathcal{D}_{\lambda_1}} = g(X,X)\xi$ for any $X \in \Gamma(\mathcal{D}_{\lambda_0})$ where ξ is the mean curvature normal of $\mathcal{D}_{\lambda_0}$. Since the distribution $\mathcal{D}_{\lambda_1}$ is totally geodesic we also have $\nabla_X X_{|\mathcal{D}_{\lambda_0}} = 0$ for any $X \in \Gamma(\mathcal{D}_{\lambda_1})$. Let $\{E_1,E_2,E_3,E_4,...,E_{2n-1},E_{2n}\}$ be a local orthonormal basis of TM such that $\mathcal{D}_{\lambda_0} = \operatorname{span}\{E_1\}$ and $\mathcal{D}_{\lambda_1} = \operatorname{span}\{E_2,E_3,E_4,...,E_{2n}\}$. Then $\nabla_{E_i}E_{i|\mathcal{D}_{\lambda_0}} = 0$ for $i \in \{2,3,4,...,2n\}$ and

$$\nabla_{E_1}E_{1|\mathcal{D}_{\lambda_1}} = \xi.$$

Consequently (note that $\nabla\lambda'_{0|\mathcal{D}_{\lambda_0}} = 0$ if and only if $\nabla\tau_{|\mathcal{D}_{\lambda_0}} = 0$),

$$\begin{aligned} \operatorname{tr}_g \nabla S = \sum_{i=1}^{n} \nabla S(E_i,E_i) &= -(S-\lambda_0\,\mathrm{id})(\nabla_{E_1}E_1) + \nabla\lambda_{0|\mathcal{D}_{\lambda_0}} \\ &= -(\lambda_1-\lambda_0)\xi \end{aligned} \tag{2.4}$$

if we assume that $\nabla\tau_{|\mathcal{D}_{\lambda_0}} = 0$. On the other hand, $\mathrm{tr}_g \nabla S_0 = \frac{\nabla\tau}{2}$ and $\mathrm{tr}_g \nabla S = \mathrm{tr}_g \nabla S_0 - \frac{\nabla\tau}{n+1}$. Consequently,

$$\mathrm{tr}_g \nabla S = \frac{(n-1)\nabla\tau}{2(n+1)} = -\frac{1}{2}\nabla\lambda_0'. \tag{2.5}$$

Thus $\xi = -\frac{1}{2(\lambda_0-\lambda_1)}\nabla\lambda_0'$. From the Lemma it follows that (M,g) is an $\mathcal{A}\oplus\mathcal{C}^{\perp}$-manifold if $2\lambda_0 + (n-4)\lambda_1$ is constant and $\nabla\tau \in \Gamma(\mathcal{D}_{\lambda_1})$. These conditions are also necessary since $\nabla\lambda_1' = 0$ if (M,g) is an $\mathcal{A}\oplus\mathcal{C}^{\perp}$-manifold and $\mathcal{D}_{\lambda_1}$ is totally geodesic. Analogously $\xi = -\frac{1}{2(\lambda_0'-\lambda_1')}\nabla\lambda_0'$ and $\nabla\lambda_0' = -\frac{n-1}{(n+1)}\nabla\tau \in \Gamma(\mathcal{D}_{\lambda_1})$, where ξ is the mean curvature normal of the umbilical distribution $\mathcal{D}_{\lambda_0}$, if (M,g) is an $\mathcal{A}\oplus\mathcal{C}^{\perp}$-manifold.$\diamondsuit$

3. Conformally Kähler Einstein-Weyl manifolds. Let g be the standard metric of $(M,[g])$. Now let us recall that $\rho^D(X,Y) = \lambda_0 g(X,Y) + \frac{n}{4}d\omega(X,Y)$. Let us assume that (M,J) is complex and $[g]$ is Hermitian i.e. $g(JX,JY) = g(X,Y)$. It follows that ρ^D is J-invariant if and only if $d\omega$ is a $(1,1)$ form, $d\omega(JX,JY) = d\omega(X,Y)$. Since $\omega(X) = g(\xi,X)$ it follows that $d\omega$ is a $(1,1)$ form iff $\nabla_{JX}\xi = J\nabla_X\xi$.

Proposition 3.1. *Let (M,J) be a compact complex manifold with conformal Hermitian structure $[g]$. Let us assume that $[g]$ is conformally Kähler and f^2g is a Kähler metric on (M,J) where g is the standard metric and $f \in C^\infty(M)$. If $(M,[g])$ is Einstein-Weyl with J-invariant Ricci tensor ρ^D then $J\xi$ is colinear with ∇f in $U = \{x : \xi_x \neq 0\}$ and ξ is a holomorphic Killing field on (M,f^2g,J).*

Proof. Let ∇ be a Levi-Civita connection of the standard metric g and ∇^1 be a Levi-Civita connection of the Kähler metric $g_1 = f^2g$. Note that ξ is a conformal field on (M,g_1), $L_\xi g_1 = L_\xi(f^2g) = 2\xi\ln f g_1 = \sigma g_1$. Every conformal field on a compact Kähler manifold is Killing (see [L]), hence consequently $\xi f = 0$ and $\xi \in \mathfrak{iso}(M,g_1)$. On a Kähler compact manifold every Killing vector field is holomorphic (see[Mor]). Thus $\xi \in \mathfrak{hol}(M,J)$. Note that

$$\nabla_X\xi = \nabla^1_X\xi - d\ln f(X)\xi - d\ln f(\xi)X + g_1(X,\xi)\nabla^1\ln f.$$

Thus

$$\begin{aligned}\nabla_{JX}\xi - J\nabla_X\xi = &-d\ln f(JX)\xi - d\ln f(\xi)JX + g_1(JX,\xi)\nabla^1\ln f \\ &+d\ln f(X)J\xi + d\ln f(\xi)JX - g_1(X,\xi)J\nabla^1\ln f.\end{aligned}$$

Hence $\nabla_{JX}\xi = J\nabla_X\xi$ if

$$-d\ln f(JX)\xi + g_1(JX,\xi)\nabla^1\ln f + d\ln f(X)J\xi - g_1(X,\xi)J\nabla^1\ln f = 0.$$

Put $X = \xi$ then we get $g_1(\xi,\xi)J\nabla^1\ln f = -d\ln f(J\xi)\xi$. It follows that in $U = \{x \in M : \xi_x \neq 0\}$ there exists a smooth function ϕ such that $\nabla^1 f = \phi J\xi$.$\diamondsuit$

Now our aim is to prove

Theorem 3.2 *Let us assume that $(M,[g],J)$ is a compact, conformally Kähler Einstein-Weyl manifold with Hermitian Ricci tensor ρ^D which is not conformally*

Einstein. Then the conformally equivalent Kähler manifold (M, g_1, J) *admits a holomorphic Killing field with a Kähler-Ricci potential. Thus* $M = \mathbb{P}(L \oplus \mathcal{O})$ *where* L *is a holomorphic line bundle over a compact Kähler Einstein manifold* (N, h) *of positive scalar curvature or is a complex projective space* $\mathbb{CP}^n$.

Proof. Let ρ, ρ^1 be the Ricci tensors of conformally related riemannian metrics $g, g_1 = f^2 g$. Then

$$\rho = \rho^1 + (n-2)f^{-1}\nabla^1 df + [f^{-1}\Delta^1 f - (n-1)f^{-2}g_1(\nabla^1 f, \nabla^1 f)]g_1.$$

Note that $\nabla^1 df(X,Y) = g_1(\nabla^1_X \nabla^1 f, Y) = g_1(X\phi J\xi, Y) + \phi g_1(J\nabla^1_X \xi, Y)$. Thus

$$\begin{aligned}(3.1)\, \rho(X,Y) - (n-2)fX\phi g(J\xi, Y) &= \rho^1(X,Y) + (n-2)f^{-1}\phi g_1(J\nabla^1_X\xi, Y) + \\ &[f^{-1}\Delta^1 f - (n-1)f^{-2}g_1(\nabla^1 f, \nabla^1 f)]g_1(X,Y),\end{aligned}$$

where $\Delta^1 f = tr_{g_1}\nabla^1 df$.

We shall show that ξ has zeros on M. If $\xi \neq 0$ on M then the function ϕ would be defined and smooth on the whole of M. Since M is compact it would imply that there exists a point $x_0 \in M$ such that $d\phi = 0$ at x_0. On the other hand the eigenvalues λ_0, λ_1 of the Ricci tensor ρ satisfy $\lambda_0 - \lambda_1 = Cg(\xi, \xi)$ where $C \neq 0$ is a real number. Since $\xi \neq 0$ it follows that the eigenvalues of ρ do not coincide at any point of M. In particular ρ is not J-invariant at x_0, a contradiction, since the right hand part of (3.1) is J-invariant. It implies that ξ is a holomorphic Killing vector field with zeros and thus has a potential τ (see [K]), i.e. there exists $\tau \in C^\infty(M)$ such that $\xi = J\nabla^1\tau$. Hence $df = -\phi d\tau$ and $d\phi \wedge d\tau = 0$. It implies that $d\phi = \alpha d\tau$. Thus we have

(3.2)

$$\begin{aligned}\rho(X,Y) + (n-2)f^{-1}\alpha d\tau(X)d\tau(Y) &= \rho^1(X,Y) - (n-2)f^{-1}\phi H^\tau(X,Y) - \\ &[f^{-1}\alpha Q + f^{-1}\phi\Delta^1\tau + (n-1)f^{-2}\phi^2 Q]g_1(X,Y).\end{aligned}$$

where $Q = g_1(\xi, \xi)$.

Note that the tensor $\widetilde{\rho}(X,Y) = \rho(X,Y) + (n-2)f^{-1}\alpha d\tau(X)d\tau(Y)$ is J-invariant. In particular $\widetilde{\rho}(\xi,\xi) = \lambda_0 g(\xi,\xi) = \lambda_o \frac{Q}{f^2}$. On the other hand $\widetilde{\rho}(\nabla^1\tau, \nabla^1\tau) = \lambda_1\frac{Q}{f^2} + (n-2)f^{-1}\alpha Q^2$. Hence $(\lambda_0 - \lambda_1)\frac{Q}{f^2} = (n-2)f^{-1}\alpha Q^2$. Since $\lambda_0 - \lambda_1 = -\frac{1}{4}(n-2)\frac{Q}{f^2}$ we get $\alpha = -\frac{1}{4f^3}$. Hence

$$d\phi = -\frac{1}{4f^3}d\tau = \frac{1}{4f^3}\frac{df}{\phi}, \tag{3.3}$$

and we get $8\phi d\phi = -d(\frac{1}{f^2})$. Hence $d(4\phi^2 + \frac{1}{f^2}) = 0$ and $4\phi^2 + \frac{1}{f^2} = C = const.$

Let us denote $\chi = (n-2)f^{-1}\phi, \sigma_0 = f^{-1}\alpha Q + f^{-1}\phi\Delta^1\tau + (n-1)f^{-2}\phi^2 Q$. Note also that the vector field $v = \nabla^1\tau$ is holomorphic and consequently $i_v\rho^1 = -\frac{1}{2}d\Delta^1\tau = -\frac{1}{2}dY$ where $Y = \Delta^1\tau$. From the equation

$$\begin{aligned}\widetilde{\rho}(X,Y) &= \rho^1(X,Y) - (n-2)f^{-1}\phi H^\tau(X,Y) - \\ &[f^{-1}\alpha Q + f^{-1}\phi\Delta^1\tau + (n-1)f^{-2}\phi^2 Q]g_1(X,Y).\end{aligned} \tag{3.4}$$

we get

$$\frac{\lambda_0}{f^2}d\tau = -\frac{1}{2}dY + \frac{1}{2}\chi dQ - \sigma_0 d\tau,$$

and

$$dY = \chi dQ - 2\sigma d\tau \tag{3.5}$$

where $\sigma = \sigma_0 + \frac{\lambda_0}{f^2}$. From (3.5) we obtain

$$d\chi \wedge dQ - 2d\sigma \wedge d\tau = 0 \tag{3.6}$$

Since $d\chi = \gamma d\tau$ we have $d\tau \wedge (\gamma dQ + 2d\sigma) = 0$ which implies $\gamma dQ + 2d\sigma = \kappa d\tau$ for a certain function κ.

Note that

$$d\sigma_0 = f^{-1}\alpha dQ + f^{-1}\phi dY + (n-1)f^{-2}\phi^2 dQ + h d\tau = \\ [-\frac{1}{4f^4} + (n-1)f^{-2}\phi^2]dQ + f^{-1}\phi dY + h d\tau.$$

On the other hand $\lambda_0 = -\frac{(n-4)Q}{4f^2} + \frac{2C_0}{n-2}$. Hence $d(\frac{\lambda_0}{f^2}) = -\frac{(n-4)}{4f^4}dQ + k d\tau$ and

$$d\sigma = [-\frac{1}{4f^4} + (n-1)f^{-2}\phi^2 - \frac{(n-4)}{4f^4}]dQ + f^{-1}\phi dY + l d\tau.$$

Since $d\chi = (n-2)d(f^{-1}\phi) = -\frac{(n-2)}{4f^4}(1-4\phi^2 f^2)d\tau$ we have $\gamma = -\frac{(n-2)}{4f^4}(1-4\phi^2 f^2)$ and

$$[-\frac{3n-8}{4f^4} + \frac{(3n-4)\phi^2}{4f^2}]dQ + 2f^{-1}\phi dY = m d\tau. \tag{3.7}$$

From equations (3.5) and (3.7) it follows that

$$dQ \wedge d\tau = dY \wedge d\tau = 0 \tag{3.8}$$

on a dense subset of M and hence everywhere.

Define $\mathcal{D} = span\{\xi, J\xi\}$ and let $\mathcal{D}^\perp$ be an orthogonal (with respect to g so also with respect to g_1) g_1) complement to $\mathcal{D}$. Both distribution are defined in an open dense subset $U = \{x : \xi_x \neq 0\}$. Let $\pi_{\mathcal{D}}, \pi_{\mathcal{D}^\perp}$ be orthogonal projections on $\mathcal{D}, \mathcal{D}^\perp$ respectively. Let us define $\omega_{\mathcal{D}}(X,Y) = g_1(J\pi_{\mathcal{D}}X, Y), \omega_{\mathcal{D}^\perp}(X,Y) = g_1(J\pi_{\mathcal{D}^\perp}X, Y)$. Then $\omega_{\mathcal{D}} + \omega_{\mathcal{D}^\perp} = \Omega$ where $\Omega(X,Y) = g_1(JX,Y)$ is the Kähler form of (M, g_1, J). Note that $\omega_{\mathcal{D}} = \frac{1}{Q}d\tau \wedge d^c\tau$. Since ξ is a holomorphic Killing field on (M, g_1, J) it follows that $H^\tau(JX,Y) = \frac{1}{2}dd^c\tau(X,Y)$. Since $\nabla^1_v v = -\frac{1}{2}\nabla^1 Q = cv$ it follows that $\mathcal{D}$ is an eigendistribution of both ρ^1 and $dd^c\tau$. We have (we denote the Ricci form also by ρ^1)

$$\rho^1 = \lambda\omega_{\mathcal{D}} + \omega_1 \tag{3.9a}$$

$$\frac{1}{2}dd^c\tau = \mu\omega_{\mathcal{D}} + \omega_2, \tag{3.9b}$$

where λ, μ are eigenvalues of ρ^1, H^τ corresponding to an eigendistribution $\mathcal{D}$. The eigenvalue μ satisfies an equation $\mu Q = H^\tau(\nabla^1\tau, \nabla^1\tau) = -\frac{1}{2}dQ(\nabla^1\tau) = -\frac{1}{2}\beta Q$ where $dQ = \beta d\tau$. Hence $\mu = -\frac{1}{2}\beta$ and $d\mu \wedge d\tau = 0$. From (3.4) it is clear that also $d\lambda \wedge d\tau = 0$. Now we have

$$\widetilde{\rho} = \rho^1 - \frac{1}{2}\chi dd^c\tau - \sigma_0\Omega, \tag{3.10}$$

and consequently

$$\frac{\lambda_0}{f^2}\omega_{\mathcal{D}} = \lambda\omega_{\mathcal{D}} - \chi\mu\omega_{\mathcal{D}} - \sigma_0\omega_{\mathcal{D}} \tag{3.11}$$

and

$$\frac{\lambda_1}{f^2}\omega_{\mathcal{D}^\perp} = \omega_1 - \chi\omega_2 - \sigma_0\omega_{\mathcal{D}^\perp}. \tag{3.12}$$

From (3.11) we obtain $\lambda - \mu\chi = \sigma_0 + \frac{\lambda_0}{f^2}$. Hence

$$\omega_1 - \chi\omega_2 = (\sigma_0 + \frac{\lambda_1}{f^2})\omega_{\mathcal{D}^\perp} = (\lambda - \mu\chi + \frac{\lambda_1 - \lambda_0}{f^2})\omega_{\mathcal{D}^\perp} = \sigma_1\omega_{\mathcal{D}^\perp}. \tag{3.13}$$

From (3.9) we get

$$\lambda d\omega_{\mathcal{D}} = -d\omega_1, \mu d\omega_{\mathcal{D}} = -d\omega_2.$$

Equation (3.13) implies that

$$d\omega_1 - d\chi \wedge \omega_2 - \chi d\omega_2 = d\sigma_1 \wedge \omega_{\mathcal{D}^\perp} + \sigma_1 d\omega_{\mathcal{D}^\perp},$$

thus

$$(-\lambda + \mu\chi + \sigma_1)d\omega_{\mathcal{D}} = d\chi \wedge \omega_2 + d\sigma_1 \wedge \omega_{\mathcal{D}^\perp}.$$

Note that $d\omega_{\mathcal{D}} = d(\frac{1}{Q}d\tau \wedge d^c\tau) = -\frac{1}{Q}d\tau \wedge dd^c\tau = -\frac{2}{Q}d\tau \wedge (\mu\omega_{\mathcal{D}} + \omega_2) = -\frac{2}{Q}d\tau \wedge \omega_2$ and $\lambda_1 - \lambda_0 = \frac{1}{4}(n-2)\frac{Q}{f^2}$. Let us write $d\sigma_1 = \psi d\tau$, then we obtain

$$d\tau \wedge (\frac{n-2}{4f^4}(-1 - 4f^2\phi^2)\omega_2 - \psi\omega_{\mathcal{D}^\perp}) = 0. \tag{3. 14}$$

From (3.14) it is clear that in U we have $\omega_2 = \kappa_2\omega_{\mathcal{D}^\perp}$ for a certain function $\kappa_2 \in C^\infty(U)$. Hence also $\omega_1 = \kappa_1\omega_{\mathcal{D}^\perp}$ for a certain function $\kappa_1 \in C^\infty(U)$. It follows that the function τ is a Kähler-Ricci potential in the sense of [D-M-1], [D-M-2], i.e. the distributions $\mathcal{D}, \mathcal{D}^\perp$ are eigendistribution of both ρ^1 and H^τ. The fact that the Einstein-Kähler manifold (N, h) has a positive scalar curvature is proved below. It is easy to check that also for $dimM = 4$ the manifold (N, h) has constant scalar curvature. The Einstein-Weyl structure on these manifolds is described in [W-W], [M-P-P-S]. ♢

4. Eigenvalues of the Ricci tensor. In our construction we shall follow L. Bérard Bergery (see [Ber]). Let (N, h, J) be a compact Kähler Einstein manifold, which is not Ricci flat and $\dim N = 2m$, $s \geq 0, L > 0, s \in \mathbb{Q}, L \in \mathbb{R}$, and $g : [0, L] \to$

$\mathbb{R}$ be a positive, smooth function on $[0, L]$ which is even at 0 and L, i.e. there exists an $\epsilon > 0$ and even, smooth functions $g_1, g_2 : (-\epsilon, \epsilon) \to \mathbb{R}$ such that $g(t) = g_1(t)$ for $t \in [0, \epsilon)$ and $g(t) = g_2(L - t)$ for $t \in (L - \epsilon, L]$. Let $f : (0, L) \to \mathbb{R}$ be positive on $(0, L)$, $f(0) = f(L) = 0$ and let f be odd at the points $0, L$. Let P be a circle bundle over N classified by the integral cohomology class $\frac{s}{2}c_1(N) \in H^2(N, \mathbb{R})$ if $c_1(N) \neq 0$. Let q be the unique positive integer such that $c_1(N) = q\alpha$ where $\alpha \in H^2(N, \mathbb{R})$ is an indivisible integral class. Such a q exists if N is simply connected or $\dim N = 2$. Note that every Kähler Einstein manifold with positive scalar curvature is simply connected. Then

$$s = \frac{2k}{q}; k \in \mathbb{Z}.$$

It is known that $q = n$ if $N = \mathbb{CP}^{n-1}$ (see [B], p.273). Note that $c_1(N) = \{\frac{1}{2\pi}\rho_N\} = \{\frac{\tau_N}{4m\pi}\omega_N\}$ where $\rho_N = \frac{\tau_N}{2m}\omega_N$ is the Ricci form of (N, h, J), τ_N is the scalar curvature of (N, h) and ω_N is the Kähler form of (N, h, J). We can assume that $\tau_N = \pm 4m$. In the case $c_1(N) = 0$ we shall assume that (N, h, J) is a Hodge manifold, i.e. the cohomology class $\{\frac{s}{2\pi}\omega_N\}$ is an integral class. On the bundle $p : P \to N$ there exists a connection form θ such that $d\theta = sp^*\omega_N$ where $p : P \to N$ is the bundle projection. Let us consider the manifold $U_{s,f,g} = (0, L) \times P$ with the metric

$$g = dt^2 + f(t)^2\theta^2 + g(t)^2 p^* h. \tag{4.1}$$

It is known that the metric (3.1) extends to a metric on the sphere bundle $M = P \times_{S^1} \mathbb{CP}^1$ if and only if a function g is positive and smooth on $[0, L]$, even at the points $0, L$, the function f is positive on $(0, L)$, smooth and odd at $0, L$ and additionally

$$f'(0) = 1, \qquad f'(L) = -1 \tag{4.2}$$

Then the metric (4.1) is bi-Hermitian. We shall prove this in Section 4. Note that $M = \mathbb{P}(L \oplus \mathcal{O})$ where $L = P \times_{S^1} \mathbb{C}$ with S^1 acting in a standard way on $\mathbb{C}$ and $\mathcal{O}$ is the trivial line bundle over N.

The metric (4.1) extends to a metric on $\mathbb{CP}^n$ if and only if the function g is positive and smooth on $[0, L)$, even at 0, odd at L, the function f is positive, smooth and odd at $0, L$ and additionally

$$f'(0) = 1, \qquad f'(L) = -1, \qquad g(L) = 0, \qquad g'(L) = -1. \tag{4.3}$$

Let us assume that (N, h) is a $2(n-1)$-dimensional Kähler-Einstein manifold of scalar curvature $4(n-1)\epsilon$ where $\epsilon \in \{-1, 0, 1\}$. Using the results in Sections 4 and 5 we obtain the following formulae for the eigenvalues of the Ricci tensor ρ of $(U_{s,f,g}, g_{f,g})$:

$$\lambda_0 = -2(n-1)\frac{g''}{g} - \frac{f''}{f}, \tag{4.2}$$

$$\lambda_1 = -\frac{f''}{f} + 2(n-1)\left(\frac{s^2 f^2}{4g^4} - \frac{f'g'}{fg}\right),$$

$$\lambda_2 = -\frac{g''}{g} + \left(\frac{s^2 f^2}{4g^4} - \frac{f'g'}{fg}\right) + \frac{2\epsilon}{g^2} - \frac{3s^2 f^2}{4g^4} - (2n-3)\frac{(g')^2}{g^2}.$$

We shall show that in fact $\epsilon = 1$, i.e. the scalar curvature of the Einstein manifold (N, h, J) is positive. From [J] it follows that the scalar curvature of Einstein-Weyl manifold λ_1 is nonnegative. We also have $\lambda_0 = \lambda_2$ and $\lambda_1 + C^2 f^2 = \lambda_0$ for a positive constant C. Since $f(0) = 0 = f(L)$ it follows that f attains a maximum at a point $t_0 \in (0, L)$. Then $f'(t_0) = 0$ and $f''(t_0) \leq 0$. Hence at t_0 we have

$$\lambda_1 = -\frac{f''}{f} + 2(n-1)\frac{s^2 f^2}{4g^4} > 0$$

and

$$-\frac{g''}{g} + \frac{2\epsilon}{g^2} - \frac{s^2 f^2}{2g^4} - (2n-3)\frac{(g')^2}{g^2} = -2(n-1)\frac{g''}{g} - \frac{f''}{f}$$

and hence

$$\frac{2\epsilon}{g^2} = \frac{s^2 f^2}{2g^4} + (2n-3)\frac{(g')^2}{g^2} - (2n-3)\frac{g''}{g} - \frac{f''}{f}.$$

From (4.2) it follows that at t_0

$$-2(n-1)\frac{g''}{g} = 2(n-1)\frac{s^2 f^2}{4g^4} + C^2 f^2 > 0$$

and consequently $\epsilon > 0$.

References.

[B] A. Besse 'Einstein manifolds' *Springer Verlag* Berlin Heidelberg (1987)

[Ber] L. Bérard Bergery,*Sur de nouvelles variétés riemanniennes d'Einstein*, Publ. de l'Institute E. Cartan (Nancy) **4**,(1982), 1-60.

[G] P. Gauduchon *La 1-forme de torsion d'une variete hermitienne compacte* Math. Ann. **267** (1984), 495-518.

[Gr] A. Gray *Einstein-like manifolds which are not Einstein* Geom. Dedicata **7** (1978) 259-280.

[D-T] M. Dunajski, K.P. Tod Four dimensional metrics conformal to Kähler to appear in Mathematical Proceedings of the Cambridge Philosophical Society.

[D-M-1] A. Derdziński, G. Maschler *Special Kähler-Ricci potentials on compact Kähler manifolds*, J. reine angew. Math. **593**, (2006), 73-116.

[D-M-2] A. Derdziński, G. Maschler *Local classification of conformally-Einstein Kähler metrics in higher dimension*, Proc. London Math. Soc. (3) **87**, (2003), no. 3, 779-819.

[D-M-3] A. Derdzinski , G. Maschler *A moduli curve for compact conformally-Einstein Khler manifolds*, Compos. Math. 141 (2005), no. 4,1029-1080

[J] W. Jelonek *Killing tensors and Einstein-Weyl geometry* Colloquium Math., vol.**81**, no.1, (1999), 5-19.

[K] S. Kobayashi *Transformation groups in differential geometry* Springer-Verlag 1972

[L] A. Lichnerowicz, Geometrie des groupes de transformations, Dunod, Paris, 1958.

[M] A. Moroianu Lectures on Kähler Geometry, LMSST 69, Cambrige University Press, 2007

[MPPS] B.Madsen, H. Pedersen, Y. Poon, A. Swann *Compact Einstein-Weyl manifolds with large symmetry group.* Duke Math. J. **88** (1997), 407-434.

[PS1] H. Pedersen and A. Swann *Riemannian submersions, four-manifolds and Einstein-Weyl geometry* Proc.London Math. Soc (3) **66** (1993) 381-399.

[PS2] H. Pedersen and A. Swann *Einstein-Weyl geometry, the Bach tensor and conformal scalar curvature* J. reine angew. Math.**441** (1993), 99-113.

[T] K.P. Tod *Compact 3-dimensional Einstein-Weyl structures* J. London Math. Soc.(2), 45 (1992) 341-351

[WW] J. Wang, M.Y. Wang *Einstein metrics on S^2-bundles* Math. Ann. 310, 497-526, (1998)

Institute of Mathematics
Cracow University of Technology
Warszawska 24
31-155 Krak'ow, POLAND.
e-mail: wjelon@usk.pk.edu.pl